abcam

Product datasheet

Anti-TDP43 antibody ab41881



★★★★★ 4 Abreviews 12 References

画像数 2

製品の概要

製品名 Anti-TDP43 antibody

製品の詳細 Rabbit polyclonal to TDP43

由来種 Rabbit

アプリケーション **適用あり:** WB 種交差性 交差種: Human

交差が予測される動物種: Mouse 🔷

免疫原 Synthetic peptide conjugated to KLH derived from within residues 350 to the C-terminus of

Human TARDBP.Immunogen の所有権に関して(Peptide available as ab41970.)

特記事項 The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

> Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any guestions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -

80°C. Avoid freeze / thaw cycle.

バッファー pH: 7.40

Preservative: 0.02% Sodium azide

Constituent: PBS

Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising

agent. If you would like information about the formulation of a specific lot, please contact our

scientific support team who will be happy to help.

精製度 Immunogen affinity purified

ポリモノ ポリクローナル

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab41881の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	****(1)	Use a concentration of 1 µg/ml. Detects a band of approximately 47 kDa (predicted molecular weight: 45 kDa).

ターゲット情報

機能 DNA and RNA-binding protein which regulates transcription and splicing. Involved in the

regulation of CFTR splicing. It promotes CFTR exon 9 skipping by binding to the UG repeated motifs in the polymorphic region near the 3'-splice site of this exon. The resulting aberrant splicing

is associated with pathological features typical of cystic fibrosis. May also be involved in microRNA biogenesis, apoptosis and cell division. Can repress HIV-1 transcription by binding to

the HIV-1 long terminal repeat. Stabilizes the low molecular weight neurofilament (NFL) mRNA

through a direct interaction with the 3' UTR.

組織特異性 Ubiquitously expressed. In particular, expression is high in pancreas, placenta, lung, genital tract

and spleen.

関連疾患 Defects in TARDBP are the cause of amyotrophic lateral sclerosis type 10 (ALS10)

[MIM:612069]. ALS is a neurodegenerative disorder affecting upper and lower motor neurons and resulting in fatal paralysis. Sensory abnormalities are absent. Death usually occurs within 2 to 5 years. The etiology of ALS is likely to be multifactorial, involving both genetic and environmental

factors. The disease is inherited in 5-10% of the cases.

配列類似性 Contains 2 RRM (RNA recognition motif) domains.

ドメイン The RRM domains can bind to both DNA and RNA.

翻訳後修飾 Hyperphosphorylated in hippocampus, neocortex, and spinal cord from individuals affected with

ALS and FTLDU.

Ubiquitinated in hippocampus, neocortex, and spinal cord from individuals affected with ALS and

FTLDU.

Cleaved to generate C-terminal fragments in hippocampus, neocortex, and spinal cord from


individuals affected with ALS and FTLDU.


細胞内局在 Nucleus. In patients with frontotemporal lobar degeneration and amyotrophic lateral sclerosis, it is


absent from the nucleus of affected neurons but it is the primary component of cytoplasmic

ubiquitin-positive inclusion bodies.

画像







Western blot - Anti-TDP43 antibody (ab41881)

Lane 1: Wild-type HAP1 cell lysate (40 µg)

Lane 2: TDP43 knockout HAP1 cell lysate (40 µg)

Lane 3: HeLa cell lysate (40 µg)

Lane 4: Jurkat cell lysate (40 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab41881 observed at 48 kDa. Red - loading control, ab8245, observed at 37 kDa.

ab41881 was shown to specifically react with TDP43 when TDP43 knockout samples were used. Wild-type and TDP43 knockout samples were subjected to SDS-PAGE. Ab41881 and ab8245 (loading control to GAPDH) were diluted at 1 µg/mL and 1/10000 dilution respectively and incubated overnight at 4C. Blots were developed with Goat anti-Rabbit lgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.

All lanes: Anti-TDP43 antibody (ab41881) at 1 µg/ml

Lane 1: HeLa (Human epithelial carcinoma cell line) Whole Cell Lysate

Lane 2: HeLa (Human epithelial carcinoma cell line) Nuclear Lysate

Lane 3 : Jurkat (Human T cell lymphoblast-like cell line) Whole Cell Lysate

Lane 4: Jurkat nuclear extract lysate (ab14844)

Lane 5: HepG2 (Human hepatocellular liver carcinoma cell line) Whole Cell Lysate

Lane 6: Hep G2 nuclear extract lysate (ab14660)

Lane 7: A431 (Human epithelial carcinoma cell line) Whole Cell Lysate

Lane 8: K-562 (Human) Nuclear Lysate - tumor cell line (ab29309)

Lysates/proteins at 10 µg per lane.

Secondary

All lanes: Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution

Performed under reducing conditions.

Predicted band size: 45 kDa **Observed band size:** 47 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors