abcam

Product datasheet

Anti-TMS1/ASC antibody [EPR10403] ab155970

יעלאעבע RabMAb

23 References 画像数6

製品の概要

製品名 Anti-TMS1/ASC antibody [EPR10403]

製品の詳細 Rabbit monoclonal [EPR10403] to TMS1/ASC

由来種 Rabbit

アプリケーション 適用あり: Flow Cyt (Intra), WB, ICC/IF

適用なし: IHC-P

種交差性 交差種: Human

免疫原 Synthetic peptide within Human TMS1/ASC aa 100 to the C-terminus. The exact sequence is

proprietary.

Database link: Q9ULZ3

ポジティブ・コントロール

特記事項

WB: THP-1 and U937 cell lysates. ICC/IF: HeLa and THP-1 cells. Flow Cyt (intra): THP-1 cells.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with

these species. Please contact us for more information.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at -20°C.

バッファー Preservative: 0.01% Sodium azide

Constituents: 40% Glycerol (glycerin, glycerine), 0.05% BSA, PBS

精製度 Protein A purified

モノクローナル ポリモノ

クローン名 EPR10403

アイソタイプ lgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab155970の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		1/10 - 1/100. ab172730 - Rabbit monoclonal lgG, is suitable for use as an isotype control with this antibody.
WB		1/1000 - 1/10000. Predicted molecular weight: 22 kDa.
ICC/IF		1/100 - 1/250.

追加情報 Is unsuitable for IHC-P.

ターゲット情報

機能 Promotes caspase-mediated apoptosis. This proapoptotic activity is mediated predominantly

through the activation of caspase-9. May be a component of the inflammasome, a protein complex which also includes NALP2, CARD8 and CASP1 and whose function would be the

activation of proinflammatory caspases.

組織特異性 Widely expressed at low levels. Detected in peripheral blood leukocytes, lung, small intestine,

spleen, thymus, colon and at lower levels in placenta, liver and kidney. Very low expression in skeletal muscle, heart and brain. Detected in the leukemia cell lines HL-60 and U937, but not in Jurkat T-cell lymphoma and Daudi Burkitt's lymphoma. Detected in the melanoma cell line WM35,

but not in WM793. Not detected in HeLa cervical carcinoma cells and Molt 4 lymphocytic

leukemia cells.

配列類似性 Contains 1 CARD domain.

Contains 1 DAPIN domain.


ドメイン Interacts with CIAS1/PYPAF1 and PYDC1 via the DAPIN domain.

翻訳後修飾 Phosphorylated.

細胞内局在 Cytoplasm. Upstream of caspase activation, a redistribution from the cytoplasm to the

aggregates occurs. These appear as hollow, perinuclear spherical, ball-like structures.

画像



All lanes : Anti-TMS1/ASC antibody [EPR10403] (ab155970) at 1/5000 dilution

Lane 1 : THP-1 (Human monocytic leukemia cell line) whole cell


Lane 2 : U937 (Human histiocytic lymphoma cell line) whole cell lysate

Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit lgG H&L (HRP) ($\underline{ab97051}$) at 20000 μg

Predicted band size: 22 kDa




Blocking and diluting buffer: 5% NFDM/TBST

All lanes : Anti-TMS1/ASC antibody [EPR10403] (ab155970) at 1/1000 dilution

Lane 1 : THP1 cell lysate Lane 2 : U937 cell lysate

Lysates/proteins at 10 µg per lane.

Predicted band size: 22 kDa




Immunocytochemistry/ Immunofluorescence - Anti-TMS1/ASC antibody [EPR10403] (ab155970)

ab155970 staining TMS1/ASC in U937 (Human histiocytic lymphoma cell line) cells by ICC/IF

(Immunocytochemistry/immunofluorescence).

Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. Samples were incubated with primary antibody at a dilution of 1/250. A goat anti rabbit IgG (Alexa Fluor[®] 488) (ab150077) was used as the secondary antibody at a dilution of 1/1000. ab195889 was used as a counterstain for primary antibody ab155970 at 1/200. DAPI was used as a nuclear counterstain and PBS as a negative control.




Flow Cytometry (Intracellular) - Anti-TMS1/ASC antibody [EPR10403] (ab155970)

ab155970 staining TMS1/ASC in the human cell line THP-1 (Human monocytic leukemia cell line) by intracellular flow cytometry.


Cells were fixed with 4% paraformaldehyde and the sample was incubated with the primary antibody at a dilution of 1/50. A goat anti rabbit lgG (Alexa Fluor[®] 488) at a dilution of 1/2000 was used as the secondary antibody.

Isoytype control: Rabbit monoclonal IgG (Black)

Unlabeled control: Cell without incubation with primary antibody and secondary antibody (Blue)



Immunocytochemistry/ Immunofluorescence - Anti-TMS1/ASC antibody [EPR10403] (ab155970) Immunofluorescence analysis of THP-1 (Human monocytic leukemia cell line) cells, labeling TMS1/ASC using ab155970 at a 1/100 dilution.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors