

### Anti-TNFAIP3 antibody [59A426] ab13597

KO 評価済

★★★★☆ 1 Abreviews 35 References 画像数 5

#### 製品の概要

|              |   |
|--------------|---|
| 製品名          | Anti-TNFAIP3 antibody [59A426]  |
| 製品の詳細        | Mouse monoclonal [59A426] to TNFAIP3  |
| 由来種          | Mouse   |
| アプリケーション     | <b>適用あり:</b> Flow Cyt (Intra), IHC-P, WB<br><b>適用なし:</b> ICC  |
| 種交差性         | <b>交差種:</b> Human   |
| 免疫原          | Recombinant full length protein corresponding to Human TNFAIP3.<br>Database link: <a href="#">P21580</a>  |
| エピトープ        | The epitope has been mapped to the C-terminal portion of A20, amino acids 440-790.  |
| ポジティブ・コントロール | WB: Daudi and HeLa cell lysates. Flow Cyt (Intra): HepG2 cells  |
| 特記事項         | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p> |

#### 製品の特性

|       |  |
|-------|--|
| 製品の状態 | Liquid   |
| 保存方法  | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle. |
| バッファー | pH: 7.40<br>Preservative: 0.02% Sodium azide<br>Constituent: PBS   |
| 精製度   | Protein G purified   |
| ポリ/モノ | モノクローナル  |
| クローン名 | 59A426   |

アイソタイプIgG1

アプリケーション

The Abpromise guarantee      **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab13597の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


| アプリケーション         | Abreviews | 特記事項   |
|------------------|-----------|--|
| Flow Cyt (Intra) |           | Use 1-2µg for 10 <sup>6</sup> cells.<br><b>ab170190</b> - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody. |
| IHC-P            |           | Use at an assay dependent concentration.   |
| WB               | ★★★★★ (1) | Use a concentration of 2 - 4 µg/ml. Detects a band of approximately 70 kDa.  |

追加情報      Is unsuitable for ICC.

ターゲット情報

|       |  |
|-------|--|
| 機能    | Ubiquitin-editing enzyme that contains both ubiquitin ligase and deubiquitinase activities.<br>Essential component of a ubiquitin-editing protein complex, comprising also RNF11, ITCH and TAX1BP1, that ensures the transient nature of inflammatory signaling pathways. Upon TNF stimulation, deubiquitinates 'Lys-63'-polyubiquitin chains on RIPK1 and catalyzes the formation of 'Lys-48'-polyubiquitin chains. This leads to RIPK1 proteasomal degradation and consequently termination of the TNF- or LPS-mediated activation of NF-kappa-B. In vitro able to deubiquitinate both 'Lys-48'- and 'Lys-63' polyubiquitin chains. Inhibitor of programmed cell death. Has a role in the function of the lymphoid system. |
| 配列類似性 | Belongs to the peptidase C64 family.<br>Contains 7 A20-type zinc fingers.<br>Contains 1 OTU domain.  |
| ドメイン  | The A20-type zinc fingers mediate the ubiquitin ligase activity.<br>The OTU domain mediates the deubiquitinase activity.   |
| 細胞内局在 | Cytoplasm. Nucleus.  |

画像



Western blot - Anti-TNFAIP3 antibody [59A426]  
(ab13597)

**All lanes :** Anti-TNFAIP3 antibody [59A426] (ab13597) at 1/500 dilution

**Lane 1 :** Wild-type HeLa cell lysate

**Lane 2 :** TNFAIP3 knockout HeLa cell lysate

**Lane 3 :** Jurkat cell treated with 5ng/ml PMA for 48 hours and then treated with 2µg/ml PHA for 48 hours, whole cell lysate

**Lane 4 :** Untreated Jurkat cell lysate


Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

**Observed band size:** 80 kDa


**Lanes 1-4:** Merged signal (red and green). Green - ab13597 observed at 80 kDa. Red - loading control, **ab181602** observed at 37 kDa.

ab13597 Anti-TNFAIP3 antibody [59A426] was shown to specifically react with TNFAIP3 in wild-type HeLa cells. Loss of signal was observed when knockout cell line **ab265983** (knockout cell lysate **ab257112**) was used. Wild-type and TNFAIP3 knockout samples were subjected to SDS-PAGE. ab13597 and Anti-GAPDH antibody [EPR16891] - Loading Control (**ab181602**) were incubated overnight at 4°C at 1 in 500 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Mouse IgG H&L (IRDye® 800CW) preadsorbed (**ab216772**) and Goat anti-Rabbit IgG H&L (IRDye® 680RD) preadsorbed (**ab216777**) secondary antibodies at 1 in 10000 dilution for 1 hour at room temperature before imaging.




Western blot - Anti-TNFAIP3 antibody [59A426]  
(ab13597)

Anti-TNFAIP3 antibody [59A426] (ab13597) at 4 µg/ml + Jurkat cell lysate



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TNFAIP3 antibody [59A426] (ab13597)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human placenta tissue labelling TNFAIP3 with ab13597 at 5µg/ml. Staining was enhanced by boiling tissue sections in 10mM sodium citrate buffer, pH6.0 for 10-20 minutes followed by cooling at room temperature for 20 minutes.



Western blot - Anti-TNFAIP3 antibody [59A426]  
(ab13597)

**All lanes** : Anti-TNFAIP3 antibody [59A426] (ab13597) at 1 µg/ml

**Lane 1** : Jurkat (Human T cell lymphoblast-like cell line) Whole Cell Lysate

**Lane 2** : HeLa (Human epithelial carcinoma cell line) Whole Cell Lysate

Lysates/proteins at 10 µg per lane.

### Secondary

**All lanes** : Goat Anti-Mouse IgG H&L (HRP) preadsorbed ([ab97040](#)) at 1/5000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

**Observed band size:** 90 kDa

**Additional bands at:** 15 kDa, 34 kDa. We are unsure as to the identity of these extra bands.

**Exposure time:** 20 minutes



Flow Cytometry (Intracellular) - Anti-TNFAIP3 antibody [59A426] (ab13597)

Overlay histogram showing HepG2 cells stained with ab13597 (red line). The cells were fixed with methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab13597, 2µg/1x10<sup>6</sup> cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) ([ab96879](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] ([ab91353](#), 2µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a decreased signal in HepG2 cells fixed with methanol (5 min)/permeabilized in 0.1% PBS-Tween used under the same conditions.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors