


Anti-TRIM5 alpha antibody ab109709

★★★★☆ [3 Abreviews](#) [1 References](#) [画像数 2](#)

製品の概要

製品名	Anti-TRIM5 alpha antibody
製品の詳細	Rabbit polyclonal to TRIM5 alpha
由来種	Rabbit
アプリケーション	適用あり: WB, IHC-P
種交差性	交差種: Human 交差が予測される動物種: Chimpanzee, Macaque monkey, Gorilla 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	This antibody gave a positive signal in the following tissue lysates: Human Placenta; Human Thymus; Human Tonsil.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab109709の使用に適用されず**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (1)	Use a concentration of 1 µg/ml. Detects a band of approximately 56 kDa (predicted molecular weight: 56 kDa).
IHC-P		Use a concentration of 1 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

ターゲット情報

機能	Isoform Alpha is a retrovirus restriction factor, which mediates species-specific, early block to retrovirus infection. Targets retroviral capsid soon after entry into the cell, and prevents reverse transcription of the virus RNA genome. Isoform Alpha trimers may make multiple contacts with the hexameric lattice of CA proteins which constitute the surface of retrovirion core, and somehow inactivate the virus. Restricts efficiently infection by N-MLV, but not HIV-1. May have E3 ubiquitin-protein ligase activity.
パスウェイ	Protein modification; protein ubiquitination.
配列類似性	Belongs to the TRIM/RBCC family. Contains 1 B box-type zinc finger. Contains 1 B30.2/SPRY domain. Contains 1 RING-type zinc finger.
ドメイン	The RING-type zinc finger domain mediates binding to an E2 ubiquitin-conjugating enzyme.
翻訳後修飾	Ubiquitinates itself in a RING finger- and UBE2D2-dependent manner (in vitro).
細胞内局在	Cytoplasm > P-body. Cytoplasmic bodies.

画像



Western blot - Anti-TRIM5 alpha antibody
(ab109709)

All lanes : Anti-TRIM5 alpha antibody (ab109709) at 1 µg/ml

Lane 1 : Human placenta tissue lysate - total protein ([ab29745](#))

Lane 2 : Tonsil (Human) Tissue Lysate

Lane 3 : Human thymus tissue lysate - total protein ([ab30146](#))

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) preadsorbed
([ab97080](#)) at 1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.


Predicted band size: 56 kDa

Observed band size: 56 kDa

Additional bands at: 31 kDa (possible isoform), 46 kDa (possible isoform)

Exposure time: 2 minutes

This antibody was raised against an immunogen that is predicted to recognize isoforms Alpha, Beta and Epsilon of TRIM5. The predicted molecular weights of isoforms Alpha, Beta and Epsilon are 56, 46 and 31 kDa respectively.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TRIM5 alpha antibody (ab109709)

IHC image of TRIM5 alpha staining in human tonsil formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab109709, 1µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors