


Anti-Urokinase antibody [EPR6273] ab133563

リコンビナント RabMAb[®]


7 References [画像数 6](#)

製品の概要

製品名	Anti-Urokinase antibody [EPR6273]
製品の詳細	Rabbit monoclonal [EPR6273] to Urokinase
由来種	Rabbit
アプリケーション	適用あり: WB, IHC-P 適用なし: Flow Cyt or ICC/IF
種交差性	交差種: Human 交差が予測される動物種: Mouse, Rat 
免疫原	Synthetic peptide within Human Urokinase aa 50-150. The exact sequence is proprietary. Database link: P00749
ポジティブ・コントロール	WB: PC-3, LnCaP and MCF7 cell lysates. IHC-P: Human kidney tissue.
特記事項	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.
解離定数 (K_D 値)	K _D = 5.19 x 10 ⁻¹¹ M



[Learn more about K_D](#)

バッファー	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol, 0.05% BSA, 59% PBS
精製度	Protein A purified
ポリモノ	モノクローナル
クローン名	EPR6273
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab133563の使用に適用されず**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
WB		1/1000. Predicted molecular weight: 49 kDa. For unpurified use at 1/100.
IHC-P		1/150. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. See IHC antigen retrieval protocols .

追加情報 Is unsuitable for Flow Cyt or ICC/IF.

ターゲット情報

機能	Specifically cleave the zymogen plasminogen to form the active enzyme plasmin.
組織特異性	Expressed in the prostate gland and prostate cancers.
配列類似性	Belongs to the peptidase S1 family. Contains 1 EGF-like domain. Contains 1 kringle domain. Contains 1 peptidase S1 domain.
翻訳後修飾	Phosphorylation of Ser-158 and Ser-323 abolishes proadhesive ability but does not interfere with receptor binding.
細胞内局在	Secreted.

画像



Western blot - Anti-Urokinase antibody [EPR6273] (ab133563)

All lanes : Anti-Urokinase antibody [EPR6273] (ab133563) at 1/100 dilution (unpurified)

Lane 1 : LnCap cell lysate

Lane 2 : PC-3 cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Peroxidase-conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 49 kDa

Observed band size: 49 kDa


Blocking buffer and concentration: 5% NFDm/TBST.

Diluting buffer and concentration: 5% NFDm /TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Urokinase antibody [EPR6273] (ab133563)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human kidney tissue labelling Urokinase with purified ab133563 at 1/150. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. A prediluted HRP-polymer conjugated anti-rabbit IgG was used as the secondary antibody. Negative control using PBS instead of primary antibody. Counterstained with Hematoxylin.



Western blot - Anti-Urokinase antibody [EPR6273] (ab133563)

All lanes : Anti-Urokinase antibody [EPR6273] (ab133563) at 1/1300 dilution (purified)

Lane 1 : LnCaP cell lysate

Lane 2 : PC-3 cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Peroxidase-conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 49 kDa

Observed band size: 49 kDa

Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.



Western blot - Anti-Urokinase antibody [EPR6273] (ab133563)

All lanes : Anti-Urokinase antibody [EPR6273] (ab133563) at 1/1000 dilution (unpurified)

Lane 1 : PC-3 cell lysate

Lane 2 : LnCaP cell lysate

Lane 3 : MCF7 cell lysate


Lysates/proteins at 10 µg per lane.

Secondary

All lanes : HRP-conjugated goat anti-rabbit IgG at 1/2000 dilution

Developed using the ECL technique.

Predicted band size: 49 kDa



Equilibrium dissociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

OI-RD Scanning - Anti-Urokinase antibody
[EPR6273] (ab133563)

Why choose a recombinant antibody?

Research with confidence
Consistent and reproducible results

Long-term and scalable supply
Recombinant technology

Success from the first experiment
Confirmed specificity

Ethical standards compliant
Animal-free production

Anti-Urokinase antibody [EPR6273] (ab133563)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors