

### Anti-VAMP2 antibody [EPR12790] ab181869

リコンビナント **RabMAb**

★★★★★ **1 Abreviews** **13 References** 画像数 13

#### 製品の概要

製品名	Anti-VAMP2 antibody [EPR12790]
製品の詳細	Rabbit monoclonal [EPR12790] to VAMP2
由来種	Rabbit
アプリケーション	<b>適用あり:</b> Flow Cyt (Intra), WB, ICC/IF, IP
種交差性	<b>交差種:</b> Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: Human fetal brain and Human cerebellum lysates ICC/IF: SH-SY5Y cells. Flow Cyt (intra): Jurkat cells.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.01% Sodium azide Constituents: 40% Glycerol, 0.05% BSA, 59% PBS
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR12790
アイソタイプ	IgG

## アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab181869の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		1/100. For unpurified use at 1/150. <b>ab172730</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB	★★★★★ (1)	1/10000 - 1/50000. Detects a band of approximately 18 kDa (predicted molecular weight: 13 kDa).
ICC/IF		1/250. <b>For unpurified use at 1/500.</b>
IP		1/150. <b>For unpurified use at 1/70.</b>

## ターゲット情報

機能	Involved in the targeting and/or fusion of transport vesicles to their target membrane.
組織特異性	Nervous system and skeletal muscle.
配列類似性	Belongs to the synaptobrevin family. Contains 1 v-SNARE coiled-coil homology domain.
細胞内局在	Cytoplasmic vesicle > secretory vesicle > synaptic vesicle membrane. Cell junction > synapse > synaptosome. Neuronal synaptic vesicles.

## 画像



Anti-VAMP2 antibody [EPR12790] (ab181869) at 1/10000 dilution (purified) + Rat heart lysate at 10 µg


### Secondary

Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/1000 dilution

**Predicted band size: 13 kDa**

Blocking buffer and concentration: 5% NFDM/TBST.


Diluting buffer and concentration: 5% NFDM /TBST.



Immunocytochemistry/ Immunofluorescence - Anti-VAMP2 antibody [EPR12790] (ab181869)

ab181869 staining VAMP2 in U87-MG (human glioblastoma) cells by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with 100% methanol. Samples were incubated with primary antibody at a dilution of 1/500. A goat anti rabbit IgG (Alexa Fluor® 488) ([ab150077](#)) was used as the secondary antibody at a dilution of 1/1000. DAPI was used as a nuclear counterstain.


**Negative control 1:** PBS only.



Immunocytochemistry/ Immunofluorescence - Anti-VAMP2 antibody [EPR12790] (ab181869)


Immunofluorescent analysis of U87-MG cells

(4% paraformaldehyde-fixed) labeling VAMP2 with ab181869 at 1/500 dilution followed by Goat anti rabbit IgG (Alexa Fluor®488) at 1/200 dilution and counterstained with Dapi.



Flow Cytometry (Intracellular) - Anti-VAMP2 antibody [EPR12790] (ab181869)

Intracellular Flow Cytometry analysis of Jurkat cells fixed with 2% paraformaldehyde labeling VAMP2 with unpurified ab181869 at 1/150 dilution followed by Goat anti rabbit IgG (FITC) at 1/150 dilution. Rabbit monoclonal IgG was used as an isotype control.



Western blot - Anti-VAMP2 antibody [EPR12790]  
(ab181869)

Anti-VAMP2 antibody [EPR12790] (ab181869) at 1/50000 dilution  
(purified) + Mouse brain lysate at 10 µg


#### Secondary

Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/100000 dilution  
(Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated)

**Predicted band size:** 13 kDa

Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.



Western blot - Anti-VAMP2 antibody [EPR12790]  
(ab181869)

Anti-VAMP2 antibody [EPR12790] (ab181869) at 1/50000 dilution  
(purified) + Human cerebellum lysate at 10 µg


#### Secondary

Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at  
1/10000 dilution

**Predicted band size:** 13 kDa

Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.



Western blot - Anti-VAMP2 antibody [EPR12790]  
(ab181869)

**All lanes :** Anti-VAMP2 antibody [EPR12790] (ab181869) at  
1/50000 dilution (unpurified)

**Lane 1 :** Human fetal brain lysate


**Lane 2 :** Human cerebellum lysate

Lysates/proteins at 10 µg per lane.

#### Secondary

**All lanes :** Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at  
1/1000 dilution

**Predicted band size:** 13 kDa




Immunoprecipitation - Anti-VAMP2 antibody  
[EPR12790] (ab181869)

ab181869 (purified) at 1/150 immunoprecipitating VAMP2 in Human cerebellum whole cell lysate. 10 ug of cell lysate was present in the input. For western blotting, a HRP-conjugated Veriblot for IP Detection Reagent (**ab131366**) (1/1,500) was used for detection. A rabbit monoclonal IgG (**ab172730**) was used instead of **ab128913** as a negative control (Lane 3).

Blocking buffer and concentration: 5% NFDM/TBST.


Diluting buffer and concentration: 5% NFDM /TBST.



Immunoprecipitation - Anti-VAMP2 antibody  
[EPR12790] (ab181869)


Western blot analysis of immunoprecipitation pellet from Human cerebellum lysate immunoprecipitated using ab181869 at 1/70 dilution.

Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at 1/1000 dilution.



Flow Cytometry (Intracellular) - Anti-VAMP2  
antibody [EPR12790] (ab181869)

Intracellular Flow Cytometry analysis of SH-SY5Y cells labelling VAMP2 with purified ab181869 at 1/100 (red). Cells were fixed with 4% paraformaldehyde. A FITC-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabelled control, cells without incubation with primary and secondary antibodies.




Immunocytochemistry/ Immunofluorescence - Anti-VAMP2 antibody [EPR12790] (ab181869)

Immunocytochemistry/Immunofluorescence analysis of U87-MG (human glioblastoma) cells labelling VAMP2 with purified ab181869 at 1/250. Cells were fixed with 100% methanol and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. **ab7291**, a mouse anti-tubulin (1/1000) and **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/1000) were also used.

Control 1: primary antibody (1/100) and secondary antibody, **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500).





Control 2: **ab7291** (1/1000) and secondary antibody, **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500).



Immunocytochemistry/ Immunofluorescence - Anti-VAMP2 antibody [EPR12790] (ab181869)

Immunofluorescent analysis of SH-SY5Y cells (human cell line) (4% paraformaldehyde-fixed) labeling VAMP2 with unpurified ab181869 at 1/500 dilution followed by Goat anti rabbit IgG (Alexa Fluor®488) at 1/200 dilution and counterstained with DAPI.

Why choose a recombinant antibody?

 <p><b>Research with confidence</b> Consistent and reproducible results</p>	 <p><b>Long-term and scalable supply</b> Recombinant technology</p>
 <p><b>Success from the first experiment</b> Confirmed specificity</p>	 <p><b>Ethical standards compliant</b> Animal-free production</p>

Anti-VAMP2 antibody [EPR12790] (ab181869)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors