

Anti-VE Cadherin antibody [EPR18229] - Intercellular Junction Marker ab205336

リコンビナント RabMAb

★★★★★ **1 Abreviews** **32 References** **画像数 6**

製品の概要

製品名	Anti-VE Cadherin antibody [EPR18229] - Intercellular Junction Marker
製品の詳細	Rabbit monoclonal [EPR18229] to VE Cadherin - Intercellular Junction Marker
由来種	Rabbit
アプリケーション	適用あり: WB, ICC/IF, IP
種交差性	交差種: Mouse
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: Mouse lung, placenta, heart, kidney and spleen lysates; bEnd.3 whole cell lysate. ICC/IF: bEnd.3 cells. IP: Mouse lung whole cell lysate; bEnd.3 whole cell lysate.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR18229

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab205336の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/1000. Detects a band of approximately 125, 90 kDa (predicted molecular weight: 88 kDa).
ICC/IF	★★★★★ (1)	1/1000.
IP		1/80.

ターゲット情報

機能	Cadherins are calcium dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. This cadherin may play a important role in endothelial cell biology through control of the cohesion and organization of the intercellular junctions. It associates with alpha-catenin forming a link to the cytoskeleton.
組織特異性	Endothelial tissues and brain.
配列類似性	Contains 5 cadherin domains.
翻訳後修飾	Phosphorylated on tyrosine residues by KDR/VEGFR-2. Dephosphorylated by PTPRB.
細胞内局在	Cell junction. Cell membrane. Found at cell-cell boundaries and probably at cell-matrix boundaries.

画像



Immunocytochemistry/ Immunofluorescence - Anti-VE Cadherin antibody [EPR18229] - Intercellular Junction Marker (ab205336)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized bEnd.3 (Mouse brain microvascular endothelial cell line) cells labeling VE Cadherin with ab205336 at 1/1000 dilution, followed by Goat anti-rabbit IgG H&L (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/500 dilution (green).

Confocal image showing membrane staining on bEnd.3 cell line.


The nuclear counterstain is DAPI (blue).

Tubulin is detected with Anti-alpha Tubulin antibody -Loading control (**ab7291**) at 1/1000 dilution and Goat Anti-Mouse IgG H&L (AlexaFluor®594) preadsorbed (**ab150120**) at 1/500 dilution (red).

The negative controls are as follows:-

-ve control 1: ab205336 at 1/1000 dilution followed by **ab150120** at 1/500 dilution.

-ve control 2: **ab7291** at 1/1000 dilution followed by **ab150077** at 1/500 dilution.



Western blot - Anti-VE Cadherin antibody [EPR18229] - Intercellular Junction Marker (ab205336)

All lanes : Anti-VE Cadherin antibody [EPR18229] - Intercellular Junction Marker (ab205336) at 1/1000 dilution

Lane 1 : Mouse lung lysate

Lane 2 : Mouse placenta lysate

Lane 3 : bEnd.3 (Mouse brain microvascular endothelial cell line) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/50000 dilution

Predicted band size: 88 kDa


Observed band size: 125,90 kDa

Exposure time: 1 second

Blocking/Dilution buffer: 5% NFDm/TBST.

Due to a high degree of glycosylation and phosphorylation, the observed MW is higher than the predicted MW. The 90kDa fragment represents the extracellular domain where the immunogen

is located.



Western blot - Anti-VE Cadherin antibody [EPR18229] - Intercellular Junction Marker (ab205336)

All lanes : Anti-VE Cadherin antibody [EPR18229] - Intercellular Junction Marker (ab205336) at 1/1000 dilution

Lane 1 : Mouse heart lysate

Lane 2 : Mouse kidney lysate

Lane 3 : Mouse spleen lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/50000 dilution


Predicted band size: 88 kDa

Observed band size: 120,90 kDa

Exposure time: 1 minute

Blocking/Dilution buffer: 5% NFDm/TBST.

Due to a high degree of glycosylation and phosphorylation, the observed MW is higher than the predicted MW. The 90kDa fragment represents the extracellular domain where the immunogen is located.



Immunoprecipitation - Anti-VE Cadherin antibody [EPR18229] - Intercellular Junction Marker (ab205336)

VE Cadherin was immunoprecipitated from 1mg of Mouse lung whole cell lysate with ab205336 at 1/80 dilution.

Western blot was performed from the immunoprecipitate using ab205336 at 1/1000 dilution.

Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG, was used as secondary antibody at 1/1500.

Lane 1: Mouse lung whole cell lysate 10ug (Input).


Lane 2: ab205336 IP in Mouse lung whole cell lysate.

Lane 3: Rabbit monoclonal IgG ([ab172730](#)) instead of ab205336 in Mouse lung whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDm/TBST.

Exposure time: 1 second.

Due to a high degree of glycosylation and phosphorylation, the observed MW is higher than the predicted MW. The 90kDa fragment represents the extracellular domain where the immunogen is located.



Immunoprecipitation - Anti-VE Cadherin antibody [EPR18229] - Intercellular Junction Marker (ab205336)

VE Cadherin was immunoprecipitated from 1mg of bEnd.3 (Mouse brain microvascular endothelial cell line) whole cell lysate with ab205336 at 1/80 dilution.

Western blot was performed from the immunoprecipitate using ab205336 at 1/1000 dilution.

Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG, was used as secondary antibody at 1/1500.

Lane 1: bEnd.3 whole cell lysate 10ug (Input).





Lane 2: ab205336 IP in bEnd.3 whole cell lysate.

Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of ab205336 in bEnd.3 whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDm/TBST.

Exposure time: 5 seconds.

Why choose a recombinant antibody?

 <p>Research with confidence Consistent and reproducible results</p>	 <p>Long-term and scalable supply Recombinant technology</p>
 <p>Success from the first experiment Confirmed specificity</p>	 <p>Ethical standards compliant Animal-free production</p>

Anti-VE Cadherin antibody [EPR18229] - Intercellular Junction Marker (ab205336)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors