


# Anti-VE Cadherin antibody - Intercellular Junction Marker ab33168

★★★★★ **40 Abreviews**   **312 References**   画像数 9

### 製品の概要

製品名	Anti-VE Cadherin antibody - Intercellular Junction Marker
製品の詳細	Rabbit polyclonal to VE Cadherin - Intercellular Junction Marker
由来種	Rabbit
アプリケーション	<b>適用あり:</b> ICC/IF, WB
種交差性	<b>交差種:</b> Mouse, Human <b>交差が予測される動物種:</b> Chicken, Cow, Pig 
免疫原	Synthetic peptide corresponding to Human VE Cadherin aa 750 to the C-terminus conjugated to keyhole limpet haemocyanin. (Peptide available as <b>ab27462</b> )
ポジティブ・コントロール	ICC/IF: HUVEC cells. WB: HUVEC cell lysate and Mouse lung tissue lysate.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS  1x PBS Batches which are <1mg/ml will contain 1% BSA, batches at 1mg/ml will not.
精製度	Immunogen affinity purified

ポリモノ                      ポリクローナル  
アイソタイプ                      IgG

## アプリケーション


**The Abpromise guarantee**                      **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab33168の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF	★★★★★ (18)	Use a concentration of 0.1 - 1 µg/ml. Abcam recommends using this product with confluent cells.
WB	★★★★★ (9)	Use a concentration of 1 µg/ml. Detects a band of approximately 115,117,120 kDa (predicted molecular weight: 88 kDa). Abcam recommends using BSA blocking with this product. Milk blocking will give a greatly reduced signal strength in WB.

## ターゲット情報


機能	Cadherins are calcium dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. This cadherin may play a important role in endothelial cell biology through control of the cohesion and organization of the intercellular junctions. It associates with alpha-catenin forming a link to the cytoskeleton.
組織特異性	Endothelial tissues and brain.
配列類似性	Contains 5 cadherin domains.
翻訳後修飾	Phosphorylated on tyrosine residues by KDR/VEGFR-2. Dephosphorylated by PTPRB.
細胞内局在	Cell junction. Cell membrane. Found at cell-cell boundaries and probably at cell-matrix boundaries.

## 画像




Anti-VE Cadherin antibody - Intercellular Junction  
Marker (ab33168)

This image is courtesy of an Abreview submitted by Simon Shen



Immunocytochemistry/ Immunofluorescence - Anti-  
VE Cadherin antibody - Intercellular Junction Marker  
(ab33168)



Western blot - Anti-VE Cadherin antibody -  
Intercellular Junction Marker (ab33168)

ab33168 staining VE Cadherin in HUVECs. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% PBS-Tween for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab33168 at 1µg/ml and **ab7291**, Mouse monoclonal [DM1A] to alpha Tubulin - Loading Control. Cells were then incubated with **ab150081**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and **ab150120**, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 594), pre-adsorbed at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.

**All lanes** : Anti-VE Cadherin antibody - Intercellular Junction  
Marker (ab33168) at 1 µg/ml

**Lane 1** : HUVEC (Human Umbilical Vein Endothelial Cell) Whole  
Cell Lysate

**Lane 2** : Mouse lung tissue lysate

Lysates/proteins at 10 µg per lane.

### Secondary

**All lanes** : Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed  
(HRP) at 1/50000 dilution

**Predicted band size:** 88 kDa


**Observed band size:** 120 kDa

**Additional bands at:** 70 kDa (possible non-specific binding)

**Exposure time:** 1 minute

**Gel type:** MOPS

**Blocking buffer:** 2% BSA



Anti-VE Cadherin antibody - Intercellular Junction Marker (ab33168)  
at 1 µg/ml + HUVEC Cell Lysate at 10 µg

### Secondary

Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed (HRP) at  
1/50000 dilution

Developed using the ECL technique.

Performed under reducing conditions.


**Predicted band size:** 88 kDa

**Observed band size:** 120 kDa

**Exposure time:** 1 minute

This blot was produced using a 4-12% Bis-tris gel under the MOPS buffer system. The gel was run at 200V for 50 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 2% Bovine Serum Albumin before being incubated with ab33168 overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP, and visualised using ECL development solution **ab133406**.

The band we observe at 115 kDa is believed to be the glycosylated form of the protein.



Western blot - Anti-VE Cadherin antibody -  
Intercellular Junction Marker (ab33168)

Anti-VE Cadherin antibody - Intercellular Junction Marker (ab33168)  
at 1 µg/ml + HUVEC Cell Lysate at 10 µg

### Secondary

Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed (HRP) at  
1/50000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

**Predicted band size:** 88 kDa


**Observed band size:** 120 kDa

**Additional bands at:** 55 kDa (possible non-specific binding)

**Exposure time:** 1 minute

This blot was produced using a 4-12% Bis-tris gel under the MOPS buffer system. The gel was run at 200V for 50 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 2% Bovine Serum Albumin before being incubated with ab33168 overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP, and visualised using ECL development solution **ab133406**.


The band we observe at 115 kDa is believed to be the glycosylated form of the protein.



Immunocytochemistry/ Immunofluorescence - Anti-VE Cadherin antibody - Intercellular Junction Marker (ab33168)

This image is courtesy of Stephen Yarwood, Inst Mol, Cell and Sys Bio, United Kingdom


ICC/IF image of VE-Cadherin staining on HUVEC cells using ab33168. The cells were incubated with the primary antibody (ab33168) and the secondary was FITC conjugated anti-rabbit used at 1:400. The cells were incubated with only the secondary antibody as a negative control.



Immunocytochemistry/ Immunofluorescence - Anti-VE Cadherin antibody - Intercellular Junction Marker (ab33168)

This image is courtesy of Ana Kasirer-Friede, Univ California-San Diego, Dept. Of Medicine, United States

ICC/IF image of VE Cadherin stained HUVEC cells. The cells were incubated with the antibody ab33168 at 1/150 (Green). The cells were also stained with Rhodamine phalloidin (Red).



Western blot - Anti-VE Cadherin antibody - Intercellular Junction Marker (ab33168)

Anti-VE Cadherin antibody - Intercellular Junction Marker (ab33168) at 1 µg/ml + HUVEC Cell Lysate at 10 µg

### Secondary

Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed (HRP) at 1/50000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

**Predicted band size:** 88 kDa


**Observed band size:** 115,117 kDa

**Additional bands at:** 45 kDa (possible non-specific binding)

**Exposure time:** 1 minute

The observed band for Cadherin 5 has a higher molecular weight of 115kDa due to glycosylation of the protein.

The immunogen used to raise this antibody has 89% homology with Cadherin 18, 88kDa, which we believe is the additional observed band at 117kDa, again due to glycosylation of the protein.



Immunocytochemistry/ Immunofluorescence - Anti-VE Cadherin antibody - Intercellular Junction Marker (ab33168)

This image is courtesy of an Abreview submitted by Kara Shumansky

ab33168 staining VE Cadherin in the endothelial cell line from Human liver by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with Paraformaldehyde. Samples were incubated with primary antibody (1/100 in PBS + 2.5% BSA + 0.1% triton) for 1 hour at 37°C. Alexa Fluor 594 Chicken anti-Rabbit IgG (H+L) Cross-Adsorbed Secon was used as the secondary antibody at 4 µg/ml.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors