

Anti-Vinculin antibody [EPR19579] ab207440

KO 評価済 リコンビナント RabMAb

3 References 画像数 6

製品の概要

製品名	Anti-Vinculin antibody [EPR19579]
製品の詳細	Rabbit monoclonal [EPR19579] to Vinculin
由来種	Rabbit
アプリケーション	適用あり: WB
種交差性	交差種: Mouse, Rat, Human
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: K562, HeLa, HEK-293T, HUVEC, MCF7, PC-3, C6, PC-12, A431, and NIH/3T3 whole cell lysates; human testis, fetal heart and fetal kidney lysates; mouse heart, kidney and spleen lysates; rat kidney and spleen lysates.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR19579

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab207440の使用に適用されず**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
WB		1/1000. Detects a band of approximately 124 kDa (predicted molecular weight: 124 kDa).

ターゲット情報

機能	Actin filament (F-actin)-binding protein involved in cell-matrix adhesion and cell-cell adhesion. Regulates cell-surface E-cadherin expression and potentiates mechanosensing by the E-cadherin complex. May also play important roles in cell morphology and locomotion.
組織特異性	Metavinculin is muscle-specific.
関連疾患	Defects in VCL are the cause of cardiomyopathy dilated type 1W (CMD1W) [MIM:611407]. Dilated cardiomyopathy is a disorder characterized by ventricular dilation and impaired systolic function, resulting in congestive heart failure and arrhythmia. Patients are at risk of premature death. Defects in VCL are the cause of cardiomyopathy familial hypertrophic type 15 (CMH15) [MIM:613255]. It is a hereditary heart disorder characterized by ventricular hypertrophy, which is usually asymmetric and often involves the interventricular septum. The symptoms include dyspnea, syncope, collapse, palpitations, and chest pain. They can be readily provoked by exercise. The disorder has inter- and intrafamilial variability ranging from benign to malignant forms with high risk of cardiac failure and sudden cardiac death.
配列類似性	Belongs to the vinculin/alpha-catenin family.
ドメイン	Exists in at least two conformations. When in the closed, 'inactive' conformation, extensive interactions between the head and tail domains prevent detectable binding to most of its ligands. It takes on an 'active' conformation after cooperative and simultaneous binding of two different ligands. This activation involves displacement of the head-tail interactions and leads to a significant accumulation of ternary complexes. The active form then binds a number of proteins that have both signaling and structural roles that are essential for cell adhesion. The N-terminal globular head (Vh) comprises of subdomains D1-D4. The C-terminal tail (Vt) binds F-actin and cross-links actin filaments into bundles. An intramolecular interaction between Vh and Vt masks the F-actin-binding domain located in Vt. The binding of talin and alpha-actinin to the D1 subdomain of vinculin induces a helical bundle conversion of this subdomain, leading to the disruption of the intramolecular interaction and the exposure of the cryptic F-actin-binding domain of Vt. Vt inhibits actin filament barbed end elongation without affecting the critical concentration of actin assembly.
翻訳後修飾	Phosphorylated; on serines, threonines and tyrosines. Phosphorylation on Tyr-1133 in activated platelets affects head-tail interactions and cell spreading but has no effect on actin binding nor on localization to focal adhesion plaques. Aceylated; mainly by myristic acid but also small amount of palmitic acid.
細胞内局在	Cytoplasm > cytoskeleton. Cell junction > adherens junction. Cell membrane. Cytoplasmic face of

adhesion plaques. Recruitment to cell-cell junctions occurs in a myosin II-dependent manner. Interaction with CTNNB1 is necessary for its localization to the cell-cell junctions.

画像



All lanes : Anti-Vinculin antibody [EPR19579] (ab207440) at 1/1000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : Vinculin knockout HeLa cell lysate

Lane 3 : HEK-293 cell lysate

Lane 4 : Jurkat cell lysate


Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 124 kDa

Lanes 1-4: Merged signal (red and green). Green - ab207440 observed at 124 kDa. Red - loading control **ab7291** observed at 50 kDa.

ab207440 Anti-Vinculin antibody [EPR19579] was shown to specifically react with Vinculin in wild-type HeLa cells. Loss of signal was observed when knockout cell line **ab265580** (knockout cell lysate **ab257795**) was used. Wild-type and Vinculin knockout samples were subjected to SDS-PAGE. ab207440 and Anti-alpha Tubulin antibody [DM1A] - Loading Control (**ab7291**) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-Vinculin antibody [EPR19579] (ab207440)

All lanes : Anti-Vinculin antibody [EPR19579] (ab207440) at 1/1000 dilution

Lane 1 : Wild-type A-431 (Human epidermoid carcinoma cell line) whole cell lysate

Lane 2 : VCL knockout A-431 (Human epidermoid carcinoma cell line) whole cell lysate

Lane 3 : HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lane 4 : Jurkat (Human T cell leukemia cell line from peripheral blood) whole cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 124 kDa

Observed band size: 124 kDa

Lanes 1 -4: Merged signal (red and green). Green - ab207440 observed at 124 kDa. Red - loading control, **ab8245** (Mouse anti-GAPDH antibody [6C5]) observed at 37kDa.

ab207440 was shown to react with VCL in A431 wild-type cells in Western blot. Loss of signal was observed when VCL knockout sample was used. A431 wild-type and VCL knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3% Milk in TBS-T (0.1% Tween®) before incubation with ab207440 and **ab8245** (Mouse anti-GAPDH antibody [6C5]) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-Vinculin antibody [EPR19579] (ab207440)

All lanes : Anti-Vinculin antibody [EPR19579] (ab207440) at 1/2000 dilution

Lane 1 : K562 (Human chronic myelogenous leukemia cell line from bone marrow) whole cell lysate

Lane 2 : HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lane 3 : HUVEC (Human umbilical vein endothelial cell line) whole cell lysate

Lane 4 : MCF7 (Human breast adenocarcinoma cell line) whole cell lysate

Lane 5 : PC-3 (Human prostate adenocarcinoma cell line) whole cell lysate

Lane 6 : Human testis lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution

Predicted band size: 124 kDa

Observed band size: 124 kDa

Blocking/Dilution buffer: 5% NFDM/TBST.

Exposure time: Lane 1-4: 3 minutes; Lane 5/6: 5 seconds.



Western blot - Anti-Vinculin antibody [EPR19579]
(ab207440)

All lanes : Anti-Vinculin antibody [EPR19579] (ab207440) at 1/1000 dilution

Lane 1 : Human fetal heart lysate

Lane 2 : Human fetal kidney lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG Peroxidase Conjugate, specific to the non-reduced form of IgG at 1/10000 dilution

Predicted band size: 124 kDa

Observed band size: 124 kDa

Exposure time: 30 seconds

Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot - Anti-Vinculin antibody [EPR19579]
(ab207440)

All lanes : Anti-Vinculin antibody [EPR19579] (ab207440) at 1/1000 dilution

Lane 1 : Mouse heart lysate

Lane 2 : Mouse kidney lysate

Lane 3 : Mouse spleen lysate

Lane 4 : Rat kidney lysate

Lane 5 : Rat spleen lysate

Lane 6 : C6 (Rat glial tumor cell line) whole cell lysate

Lane 7 : PC-12 (Rat adrenal gland pheochromocytoma cell line) whole cell lysate

Lane 8 : NIH/3T3 (Mouse embryonic fibroblast cell line) whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution


Predicted band size: 124 kDa

Observed band size: 124 kDa

Blocking/Dilution buffer: 5% NFD/MTBST.

Exposure times: Lane 1-3: 30 seconds; Lane 4-8: 10 seconds.

Why choose a recombinant antibody?



- Research with confidence**
Consistent and reproducible results
- Long-term and scalable supply**
Recombinant technology
- Success from the first experiment**
Confirmed specificity
- Ethical standards compliant**
Animal-free production

Anti-Vinculin antibody [EPR19579] (ab207440)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors