

Anti-Visfatin antibody [EPR21980] ab236874

リコンビナント RabMAb

11 References [画像数 12](#)

製品の概要

製品名	Anti-Visfatin antibody [EPR21980]
製品の詳細	Rabbit monoclonal [EPR21980] to Visfatin
由来種	Rabbit
アプリケーション	適用あり: WB, IHC-P, ICC/IF, IP, Flow Cyt (Intra)
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: K562, Raji, HeLa, HT-1080, PC-3, MCF7, HCT 116, C6, PC-12 and Raw 264.7 whole cell lysate; Human fetal kidney lysate. IHC-P: Human bladder cancer and breast cancer tissue; Mouse kidney tissue; Rat liver tissue. ICC/IF: HCT 116 and K562 cells. Flow: K562 and HCT 116 cells. IP: K562 whole cell lysate.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol, 0.05% BSA
精製度	Protein A purified
ポリクローナル	モノクローナル
クローン名	EPR21980

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab236874の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/1000. Predicted molecular weight: 55 kDa.
IHC-P		1/2000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).
ICC/IF		1/100.
IP		1/30.
Flow Cyt (Intra)		1/500.

ターゲット情報

機能	Catalyzes the condensation of nicotinamide with 5-phosphoribosyl-1-pyrophosphate to yield nicotinamide mononucleotide, an intermediate in the biosynthesis of NAD. It is the rate limiting component in the mammalian NAD biosynthesis pathway.
組織特異性	Expressed in large amounts in bone marrow, liver tissue, and muscle. Also present in heart, placenta, lung, and kidney tissues.
パスウェイ	Cofactor biosynthesis; NAD(+) biosynthesis; nicotinamide D-ribonucleotide from 5-phospho-alpha-D-ribose 1-diphosphate and nicotinamide: step 1/1.
配列類似性	Belongs to the NAPRTase family.
細胞内局在	Cytoplasm.

画像



All lanes : Anti-Visfatin antibody [EPR21980] (ab236874) at 1/1000 dilution

Lane 1 : K562 (human chronic myelogenous leukemia lymphoblast), whole cell lysate at 20 µg

Lane 2 : Raji (human Burkitt's lymphoma B lymphocyte), whole cell lysate at 20 µg

Lane 3 : HeLa (human cervix adenocarcinoma epithelial cell), whole cell lysate at 20 µg

Lane 4 : HT-1080 (human fibrosarcoma epithelial cell), whole cell lysate at 20 µg

Lane 5 : PC-3 (human prostate adenocarcinoma epithelial cell), whole cell lysate at 20 µg

Lane 6 : Human fetal kidney lysate at 10 µg

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution

Predicted band size: 55 kDa

Blocking/Diluting buffer and concentration: 5% NFDM/TBST

Exposure times:


Lanes 1-2: 26 seconds

Lanes 3-5: 92 seconds

Lane 6: 3 minutes

Lane 6 was developed using a higher sensitivity ECL substrate.

The molecular weight observed is consistent with what has been described in the literature (PMID:25368373, PMID:12782293).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Visfatin antibody [EPR21980] (ab236874)

Immunohistochemical analysis of paraffin-embedded human bladder cancer tissue labeling Visfatin using ab236874 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use.

Nuclear and cytoplasmic staining in human bladder cancer (PMID:26396920) is observed. Counterstained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-Visfatin antibody [EPR21980] (ab236874)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HCT 116 (human colorectal carcinoma epithelial cell) cells labeling Visfatin (Green) with ab236874 at 1/100 dilution, followed by AlexaFluor[®]488 Goat anti-Rabbit secondary (**ab150077**) at 1/1000 dilution. Confocal image showing cytoplasmic and nuclear staining in HCT 116 cell line is observed. The nuclear counterstain is DAPI (Blue). Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor[®] 594) (**ab195889**) was used as the counterstain (Red) at 1/200 dilution.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Alexa-Fluor[®]488 Goat anti-Rabbit secondary (**ab150077**) at 1/1000 dilution.



Immunoprecipitation - Anti-Visfatin antibody
[EPR21980] (ab236874)

Visfatin was immunoprecipitated from 0.35 mg of K562 (human chronic myelogenous leukemia lymphoblast) whole cell lysate with ab236874 at 1/30 dilution. Western blot was performed from the immunoprecipitate using ab236874 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used for detection at 1/5000 dilution.


Lane 1: K562 whole cell lysate 10 µg (input).

Lane 2: ab236874 IP in K562 whole cell lysate.

Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of ab236874 in K562 whole cell lysate.


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 30 seconds.




Flow Cytometry (Intracellular) - Anti-Visfatin antibody
[EPR21980] (ab236874)

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed, 90% methanol permeabilized HCT 116 (human colorectal carcinoma epithelial cell) cell line labeling Visfatin with ab236874 at 1/500 dilution (Red) compared with the Rabbit monoclonal IgG (**ab172730**, Black) isotype control and an unlabeled control (cells without incubation with primary antibody and secondary antibody, Blue). Goat anti rabbit IgG (Alexa Fluor[®] 488, **ab150077**) at 1/2000 dilution was used as the secondary antibody.



Flow Cytometry (Intracellular) - Anti-Visfatin antibody
[EPR21980] (ab236874)

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed, 90% methanol permeabilized K562 (human chronic myelogenous leukemia lymphoblast) cell line labeling Visfatin with ab236874 at 1/500 dilution (Red) compared with the Rabbit monoclonal IgG (**ab172730**, Black) isotype control and an unlabeled control (cells without incubation with primary antibody and secondary antibody, Blue). Goat anti rabbit IgG (Alexa Fluor[®] 488, **ab150077**) at 1/2000 dilution was used as the secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Visfatin antibody [EPR21980] (ab236874)

Immunohistochemical analysis of paraffin-embedded rat liver tissue labeling Visfatin using ab236874 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Nuclear and cytoplasmic staining in rat liver (PMID: 29104634; PMID:24603648) is observed. Counterstained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Visfatin antibody [EPR21980] (ab236874)

Immunohistochemical analysis of paraffin-embedded mouse kidney tissue labeling Visfatin using ab236874 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Nuclear and cytoplasmic staining in mouse kidney (PMID: 29104634; PMID:24287278) is observed. Counterstained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Visfatin antibody [EPR21980] (ab236874)

Immunohistochemical analysis of paraffin-embedded human breast cancer tissue labeling Visfatin using ab236874 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Nuclear and cytoplasmic staining in human breast cancer (PMID:21784959) is observed. Counterstained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Western blot - Anti-Visfatin antibody [EPR21980] (ab236874)

All lanes : Anti-Visfatin antibody [EPR21980] (ab236874) at 1/1000 dilution

Lane 1 : MCF7 (human breast adenocarcinoma epithelial cell), whole cell lysate at 20 µg

Lane 2 : HCT 116 (human colorectal carcinoma epithelial cell), whole cell lysate at 20 µg

Lane 3 : HT-1080 (human fibrosarcoma epithelial cell), whole cell lysate at 20 µg

Lane 4 : C6 (rat glial tumor glial cell), whole cell lysate at 20 µg

Lane 5 : PC-12 (rat adrenal gland pheochromocytoma), whole cell lysate at 20 µg

Lane 6 : RAW264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage), whole cell lysate at 10 µg

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/100000 dilution

Predicted band size: 55 kDa

Blocking/Diluting buffer and concentration: 5% NFDm/TBST


Exposure time:

Lanes 1-3: 40 seconds

Lanes 4-5: 3 minutes

Lane 6: 15 seconds

The molecular weight observed is consistent with what has been described in the literature (PMID:25368373, PMID:12782293).



Immunocytochemistry/ Immunofluorescence - Anti-Visfatin antibody [EPR21980] (ab236874)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized K562 (human chronic myelogenous leukemia lymphoblast) cells labeling Visfatin (Green) with ab236874 at 1/100 dilution, followed by AlexaFluor[®]488 Goat anti-Rabbit secondary ([ab150077](#)) at 1/1000 dilution. Confocal image showing cytoplasmic and nuclear staining in K562 cell line is observed. The nuclear counterstain is DAPI (Blue). Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor[®] 594) ([ab195889](#)) was used as the counterstain (Red) at 1/200 dilution.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Alexa-Fluor[®]488 Goat anti-Rabbit secondary ([ab150077](#)) at 1/1000 dilution.

Why choose a recombinant antibody?

- Research with confidence**
Consistent and reproducible results
- Long-term and scalable supply**
Recombinant technology
- Success from the first experiment**
Confirmed specificity
- Ethical standards compliant**
Animal-free production

Anti-Visfatin antibody [EPR21980] (ab236874)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors