abcam

Product datasheet

Anti-ZDHHC13 antibody ab28759

2 References 画像数 1

製品の概要

免疫原

製品名 Anti-ZDHHC13 antibody

製品の詳細 Goat polyclonal to ZDHHC13

由来種 Goat

特異性 This antibody is expected to recognise both reported isoforms (as represented by NP_061901

and NP_001001483).

アプリケーション 適用あり: WB

種交差性 交差種: Human

交差が予測される動物種: Mouse, Rat 4

Synthetic peptide:

(C) FHPAREKVLRSV

, corresponding to C terminal amino acids 611-622 of Human ZDHHC13 Peptide is available as

ab45682 - ZDHHC13 peptide (611-622)

Run BLAST with
Run BLAST with

特記事項

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw

cycles.

バッファー pH: 7.30

Preservative: 0.02% Sodium azide Constituents: 0.5% Tris, 0.5% BSA

精製度 Immunogen affinity purified

特記事項(精製) This antibody was purified from goat serum by ammonium sulphate precipitation followed by

1

antigen affinity chromatography using the immunizing peptide.

ポリモノ ポリクローナル

アイソタイプ ΙgG

アプリケーション

Abpromise保証は、次のテスト済みアプリケーションにおけるab28759の使用に適用されます The Abpromise guarantee

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		

追加情報 WB: Use at a concentration of 0.1 - 3 µg/ml.

This antibody gave a positive result in ELISA against the immunizing peptide.

Antibody detection limit dilution 1:32,000.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

ターゲット情報

機能 Palmitoyltransferase for HD and GAD2 (By similarity). Mediates Mg(2+) transport.

配列類似性 Belongs to the DHHC palmitoyltransferase family. AKR/ZDHHC17 subfamily.

Contains 6 ANK repeats.


Contains 1 DHHC-type zinc finger.

ドメイン The DHHC domain is required for palmitoyltransferase activity.

細胞内局在 Golgi apparatus membrane. Cytoplasmic vesicle membrane. Low extracellular Mg(2+) induces

increase in Golgi and in post-Golgi vesicles.

画像



Anti-ZDHHC13 antibody (ab28759) at 0.1 µg/ml + Human cerebellum lysate at 35 µg

Developed using the ECL technique.

Primary antibody was incubated for 1 hour.

Western blot - Anti-ZDHHC13 antibody (ab28759)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors