

Recombinant human Caspase-9 protein ab52203

2 References [画像数 1](#)

製品の詳細

製品名	Recombinant human Caspase-9 protein
生理活性	Specific activity: 400 units/mg. Unit definition: One unit of the recombinant caspase-9 is the enzyme activity that cleaves 1 nmol of the caspase substrate LEHD-pNA (pNA: pnitroaniline) per hour at 37°C in a reaction solution containing 50 mM Hepes, pH 7.2, 50 mM NaCl, 0.1% Chaps, 10 mM EDTA, 5% Glycerol, and 10 mM DTT.
精製度	> 90 % SDS-PAGE.
発現系	Escherichia coli
アクセッション番号	<u>P55211</u>
タンパク質長	Full length protein
Animal free	No
由来	Recombinant
生物種	Human

特性

Our **Abpromise guarantee** covers the use of **ab52203** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	SDS-PAGE
製品の状態	Lyophilized
備考	This product is manufactured by BioVision, an Abcam company and was previously called 1089 Caspase-9, human recombinant. 1089-25 is the same size as the 25 unit size of ab52203. UNIT DEFINITION: One unit of the recombinant Caspase 9 is the enzyme activity that cleaves 1 nmol of the caspase substrate LEHD-pNA (pNA: pnitroaniline) per hour at 37°C in a reaction solution containing 50 mM Hepes, pH 7.2, 50 mM NaCl, 0.1% Chaps, 10 mM EDTA, 5% Glycerol, and 10 mM DTT.

前処理および保存

保存方法および安定性	Shipped at 4°C. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle.
------------	---

Constituents: PBS, 15% Glycerol (glycerin, glycerine)

This product is an active protein and may elicit a biological response in vivo, handle with caution.

再構成

Reconstitute to 1 unit per μ l in PBS containing 15% glycerol. Following reconstitution in PBS, the enzyme should be aliquoted and immediately stored at -70°C .

関連情報

機能

Involved in the activation cascade of caspases responsible for apoptosis execution. Binding of caspase-9 to Apaf-1 leads to activation of the protease which then cleaves and activates caspase-3. Proteolytically cleaves poly(ADP-ribose) polymerase (PARP). Isoform 2 lacks activity is an dominant-negative inhibitor of caspase-9.

組織特異性

Ubiquitous, with highest expression in the heart, moderate expression in liver, skeletal muscle, and pancreas. Low levels in all other tissues. Within the heart, specifically expressed in myocytes.

配列類似性

Belongs to the peptidase C14A family.
Contains 1 CARD domain.


発生段階

Expressed at low levels in fetal heart, at moderate levels in neonate heart, and at high levels in adult heart.

翻訳後修飾

Cleavages at Asp-315 by granzyme B and at Asp-330 by caspase-3 generate the two active subunits. Caspase-8 and -10 can also be involved in these processing events.

画像



Lane 1: Recombinant caspase 9 protein (Active) ab52203- 100 ng.
Lane 2: BSA- 400 ng. Lane 3: BSA- 200 ng. Lane 4: BSA- 100 ng.
Lane 5: BSA- 50 ng. Lane 6: BSA- 25 ng. SYPRO Ruby protein stain. Predicted band size: 46 kDa (precursor protein) and 35 kDa (p35 subunit) Observed Bands: 20 kDa and 11 kDa.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors