

Recombinant human Cdk4 + Cyclin D1 protein ab55695

6 References [画像数 2](#)

製品の詳細

製品名	Recombinant human Cdk4 + Cyclin D1 protein
生理活性	Specific Activity is lot specific. Please contact our technical team to get lot specific information.
精製度	> 75 % Densitometry. Affinity purified.
発現系	Baculovirus infected Sf9 cells
タンパク質長	Full length protein
Animal free	No
由来	Recombinant
生物種	Human
タグ	GST tag N-Terminus
配列の追加情報	MW: Cdk4 = 57 kDa and CyclinD1 = 61 kDa.

特性

Our **Abpromise guarantee** covers the use of **ab55695** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Western blot Functional Studies
製品の状態	Liquid
備考	14nmol/min/mg using protein substrate Rb (773-928). ab56270 (Human Rb protein fragment) can be utilized as a substrate for assessing Kinase activity

前処理および保存


保存方法および安定性	Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles. pH: 7.50 Constituents: 0.00174% PMSF, 0.00385% (R*,R*)-1,4-Dimercaptobutan-2,3-diol, 0.79% Tris HCl, 0.00292% EDTA, 25% Glycerol (glycerin, glycerine), 0.88% Sodium chloride, 0.31% Glutathione
------------	--

This product is an active protein and may elicit a biological response in vivo, handle with caution.


関連情報

関連性	CDK4 is a member of the cyclin-dependent protein kinase family and is involved in the control of cell proliferation during the G1 phase. Phosphorylation at Thr-172 is necessary for enzymatic activity. CDK4 mutations are involved in tumor formation, with defects in CDK4 resulting in cutaneous malignant melanoma 3 (CMM3). Cyclin D1 is essential for the control of the cell cycle at the G1/S transition. It interacts with the CDK4 and CDK6 protein kinases to form a serine/threonine kinase holoenzyme complex. The cyclin subunit imparts substrate specificity to the complex.
細胞内局在	Nuclear

画像



SDS-PAGE analysis of ab55695 with molecular weight markers.
Approximate molecular weights: 57kDa (CDK4)/61kDa (CyclinD1).



Kinase assay using ab55695 using protein substrate Rb. The specific activity of CDK4/CyclinD1 is 14 nmol/min/mg.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors