

Recombinant Human STAT1 alpha protein - GST tagged ab140412

[1 References](#) [画像数 1](#)

製品の詳細

製品名	Recombinant Human STAT1 alpha protein - GST tagged
精製度	> 90 % Densitometry. Affinity purified.
発現系	Baculovirus infected Sf9 cells
アクセッション番号	P42224
タンパク質長	Full length protein
Animal free	No
由来	Recombinant
生物種	Human
配列	MSQWYELQQLDSKFLEQVHQLYDDSFPMERQYLAQWLEKQD WEHAANDV SFATIRFHDLLSQLDDQYSRFSLENNFLLQHNIRKSKRNLQD NFQEDPIQ MSMIIYSCLKEERKILENAQRFNQAQSGNIQSTVMLDKQKEL DSKVRNVK DKVMCIEHEIKSLEDLQDEYDFKCKTLQNHETNGVAKSDQ KQEQLLLK KMYLMLDNKRKEVVHKIIELLNVTELTQNALINDELVEWKRR QQSACIGG PPNAQLDQLQNWFTIVAESLQVVRQQLKKLEELQKYTYEHD PITKNKQV LWDRTFSLFQQLIQSSFVVERQPCMPHPQRPLVLKTGVQFT VKLRLLVK LQELNYNLKVKVLFDKDVNERNTVKGFRKFNILGHTTKVMMN EESTNGSL AAEFRHLQLKEQKNAGTRTNEGPLIVTEELHSLSFETQLCQP GLVIDLET TSLPVVVISNVSQLPSGWASILWYNMLVAEPRNLSFFLTPPC ARWAQLSE VLSWQFSSVTKRGLNVDQLNMLGEKLLGPNASPDGLIPWTRF CKENINDK NFPFWLWIESILELIKHLLPLWNDGCIMGFISKERERALLK DQQPGTFL

LRFSESSREGAITFTWVERSQNGGEPDFHAVEPYTKKELSAV
TFPDIIRN
YKVMAAENIPENPLKYLPNIDKDHAFGKYYSRPKEAPEPME
LDGPKGTG
YIKTELISVSEVHPSRLQTTDNLLPMSPEEFDEVSRIVGSVE
FDSMMNTV

予測される分子量 110 kDa including tags
領域 1 to 750
タグ GST tag N-Terminus

特性

Our **Abpromise guarantee** covers the use of **ab140412** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション Western blot
SDS-PAGE

製品の状態 Liquid

前処理および保存

保存方法および安定性 Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles.
pH: 7.50
Constituents: 0.31% Glutathione, 0.002% PMSF, 0.004% DTT, 0.79% Tris HCl, 0.003% EDTA, 25% Glycerol (glycerin, glycerine), 0.88% Sodium chloride

関連情報

機能 Signal transducer and activator of transcription that mediates signaling by interferons (IFNs). Following type I IFN (IFN-alpha and IFN-beta) binding to cell surface receptors, Jak kinases (TYK2 and JAK1) are activated, leading to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize, associate with ISGF3G/IRF-9 to form a complex termed ISGF3 transcription factor, that enters the nucleus. ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of interferon stimulated genes, which drive the cell in an antiviral state. In response to type II IFN (IFN-gamma), STAT1 is tyrosine- and serine-phosphorylated. It then forms a homodimer termed IFN-gamma-activated factor (GAF), migrates into the nucleus and binds to the IFN gamma activated sequence (GAS) to drive the expression of the target genes, inducing a cellular antiviral state.

関連疾患 Note=STAT1 deficiency results in impaired immune response leading to severe mycobacterial and viral diseases. In the case of complete deficiency, patients can die of viral disease. Defects in STAT1 are a cause of mendelian susceptibility to mycobacterial disease (MSMD) [MIM:209950]; also known as familial disseminated atypical mycobacterial infection. This rare condition confers predisposition to illness caused by moderately virulent mycobacterial species, such as Bacillus Calmette-Guerin (BCG) vaccine and environmental non-tuberculous mycobacteria, and by the more virulent Mycobacterium tuberculosis. Other microorganisms rarely cause severe clinical disease in individuals with susceptibility to mycobacterial infections, with the exception of Salmonella which infects less than 50% of these individuals. The pathogenic

mechanism underlying MSMD is the impairment of interferon-gamma mediated immunity whose severity determines the clinical outcome. Some patients die of overwhelming mycobacterial disease with lepromatous-like lesions in early childhood, whereas others develop, later in life, disseminated but curable infections with tuberculoid granulomas. MSMD is a genetically heterogeneous disease with autosomal recessive, autosomal dominant or X-linked inheritance.

配列類似性

Belongs to the transcription factor STAT family.

Contains 1 SH2 domain.

翻訳後修飾

Phosphorylated on tyrosine and serine residues in response to IFN-alpha, IFN-gamma, PDGF and EGF. Phosphorylation on Tyr-701 (lacking in beta form) by JAK promotes dimerization and subsequent translocation to the nucleus. Phosphorylation on Ser-727 by several kinases including MAPK14, ERK1/2 and CAMKII on IFN-gamma stimulation, regulates STAT1 transcriptional activity. Phosphorylation on Ser-727 promotes sumoylation though increasing interaction with PIAS. Phosphorylation on Ser-727 by PKCdelta induces apoptosis in response to DNA-damaging agents.


Sumoylated by SUMO1, SUMO2 and SUMO3. Sumoylation is enhanced by IFN-gamma-induced phosphorylation on Ser-727, and by interaction with PIAS proteins. Enhances the transactivation activity.

ISGylated.

細胞内局在

Cytoplasm. Nucleus. Translocated into the nucleus in response to IFN-gamma-induced tyrosine phosphorylation and dimerization.

画像



SDS-PAGE analysis of ab140412.

SDS-PAGE - Recombinant Human STAT1 alpha protein (ab140412)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors