abcam

Product datasheet

Recombinant human VEGFA protein ab50078

2 References

製品の詳細

製品名 Recombinant human VEGFA protein

生理活性 Determined by its ability to stimulate the proliferation of human umbilical vein endothelial cells

(HUVEC). The expected ED $_{50}$ for this effect is 1.0-5.0 ng/ml.

精製度 > 95 % SDS-PAGE.

ab50078 purity was determined also by HPLC Endotoxin level is less than 0.1 ng per µg

(1EU/µg).

発現系 Escherichia coli

アクセッション番号 <u>P15692-9</u>

タンパク質長 Full length protein

Animal free No

由来 Recombinant

生物種 Human

配列 APMAEGGGQN HHEVVKFMDV YQRSYCHPIE

TLVDIFQEYP DEIEYIFKPS CVPLMRCGGC
CNDEGLECVP TEESNITMQI MRIKPHQGQH
IGEMSFLQHN KCECRPKKDR ARQENCDKPR R

領域 27 to 232

配列の追加情報 Recombinant Human VEGF121 is a 28.4 kDa, disulfide-linked homodimeric protein consisting of

two 121 amino acid polypeptide chains.

特性

Our Abpromise guarantee covers the use of ab50078 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション SDS-PAGE

Functional Studies

製品の状態 Lyophilized

前処理および保存

保存方法および安定性 Shipped at 4°C. The lyophilized protein is stable for a few weeks at room temperature. Store at -

1

20°C long term.

This product is an active protein and may elicit a biological response in vivo, handle with caution.

再構成 Reconsitute to 1mg/ml in distilled water

関連情報

機能 Growth factor active in angiogenesis, vasculogenesis and endothelial cell growth. Induces

endothelial cell proliferation, promotes cell migration, inhibits apoptosis and induces

permeabilization of blood vessels. Binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin. NRP1/Neuropilin-1 binds isoforms VEGF-165 and VEGF-145. Isoform VEGF165B binds to KDR but does not activate downstream signaling pathways, does

not activate angiogenesis and inhibits tumor growth.

組織特異性 Isoform VEGF189, isoform VEGF165 and isoform VEGF121 are widely expressed. Isoform

VEGF206 and isoform VEGF145 are not widely expressed.

関連疾患 Defects in VEGFA are a cause of susceptibility to microvascular complications of diabetes type 1

(MVCD1) [MIM:603933]. These are pathological conditions that develop in numerous tissues and organs as a consequence of diabetes mellitus. They include diabetic retinopathy, diabetic nephropathy leading to end-stage renal disease, and diabetic neuropathy. Diabetic retinopathy remains the major cause of new-onset blindness among diabetic adults. It is characterized by

vascular permeability and increased tissue ischemia and angiogenesis.

配列類似性 Belongs to the PDGF/VEGF growth factor family.

細胞内局在 Secreted. VEGF121 is acidic and freely secreted. VEGF165 is more basic, has heparin-binding

properties and, although a signicant proportion remains cell-associated, most is freely secreted. VEGF189 is very basic, it is cell-associated after secretion and is bound avidly by heparin and the extracellular matrix, although it may be released as a soluble form by heparin, heparinase or

plasmin.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors