

# Biotinylated Goat Anti-Rabbit IgG (H+L) (Ready to use) ab64256

★★★★★ [1 Abreviews](#) [101 References](#) [画像数 1](#)

### 製品の概要

製品名	Biotinylated Goat Anti-Rabbit IgG (H+L) (Ready to use)
アプリケーション	適用あり: IHC-P
特記事項	<p><b>Staining protocol</b></p> <p>Deparaffinize and rehydrate formalin-fixed paraffin-embedded tissue section.</p> <p>Add appropriate hydrogen peroxide blocking solution to cover the sections. Incubate for 10 minutes. Wash 2 times in buffer.</p> <p>Perform appropriate pretreatment if required. Wash 3 times in buffer.</p> <p>Apply appropriate protein blocking solution and incubate for 5 minutes at room temperature to block non specific background staining. Wash 1 time in buffer.</p> <p>Apply primary antibody and incubate according to manufacturer's protocol.</p> <p>Wash 4 times in buffer. Apply Ready to use Biotinylated Goat anti Rabbit IgG (ab64256) solution and incubate for 10-15 minutes at room temperature. Wash 4 times in buffer.</p> <p>Apply enzyme labelled streptavidin and incubate according to manufacturer's protocol.</p> <p>Rinse 4 times in buffer. Place slide in appropriate chromogenic substrate and incubate until desired reaction is achieved. Rinse 4 times in buffer.</p> <p>Add enough drops of Haematoxylin to cover the section. Incubate for 1 minute.</p> <p>Rinse 7-8 times in tap water. Add Mounting Medium to cover the section.</p>

This reagent is intended to be used in a labeled streptavidin biotin immunoenzymatic antigen detection system. This technique involves the sequential incubation of the specimen with an unconjugated primary antibody specific to the target antigen, a biotinylated secondary antibody which reacts with the primary antibody, enzyme labeled streptavidin, and substrate chromogen.

### 製品の特性

製品の状態	Liquid
保存方法	Store at +4°C.
バッファー	<p>pH: 7.2</p> <p>Preservative: 0.0985% Sodium azide</p> <p>Constituent: PBS</p>
IHC 試薬 備考	<p>This reagent is intended to be used in a labeled streptavidin biotin immunoenzymatic antigen detection system. This technique involves the sequential incubation of the specimen with an unconjugated primary antibody specific to the target antigen, a biotinylated secondary antibody</p>

which reacts with the primary antibody, enzyme labeled streptavidin, and substrate chromogen.

細胞内局在

Secreted

細胞内局在


Secreted

## アプリケーション

**The Abpromise guarantee** Abpromise保証は、 次のテスト済みアプリケーションにおけるab64256の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use at an assay dependent dilution.

## 画像



Ab64256 was used with the primary antibody [ab3517](#) in IHC-P.  
See Abreview on [ab3517](#).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Biotinylated Goat Anti-Rabbit IgG (H+L) (Ready to use) (ab64256)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors