

Immunoassay Blocking Buffer ab171534

1 References [画像数 1](#)

製品の概要

製品名	Immunoassay Blocking Buffer
アプリケーション	適用あり: ELISA 適用なし: IHC-Fr or IHC-P
特記事項	<p>Abcam's Immunoassay Blocking Buffer (ab171534) effectively preserves the conformation and activity of dried proteins in immunoassays. Immunoassay Blocking Buffer simultaneously blocks and stabilizes with superior results. It can be used to stabilize antibodies, antigens or enzymes on an assortment of immunoassay components, such as polystyrene plates, tubes, glass, membranes, and filter paper. It is easily incorporated into most assay protocols by simply substituting it for the blocking solution.</p> <p>Recommendations for Use: <i>The following are general guidelines only.</i></p> <p>To Stabilize Adsorbed or Immobilized Proteins on Microtiter Plates/Strips</p> <ol style="list-style-type: none">1. Immobilize or adsorb the primary protein (antibody or antigen) according to the procedure optimized in your laboratory; wash adequately to remove excess or weakly bound protein2. Immediately after washing, add Immunoassay Blocking Buffer stabilizer to allow interaction with the entire protein-coated surface; do not let the components dry before adding stabilizer since drying contributes to the loss of protein activity3. Incubate 15-60 minutes at room temperatures; for most assays, Immunoassay Blocking Buffer stabilizer can replace the blocking solution4. Remove or aspirate the Immunoassay Blocking Buffer stabilizer, but do not wash5. Dry components for long-term storage. Wells coated with Immunoassay Blocking Buffer stabilizer may require longer drying times than those without; drying times should be optimized for each application. Recommended methods are:<ul style="list-style-type: none">- Place plates in a humidity-controlled chamber until dry (4-24 hours)- Dry plates at 30°C-40°C in a vacuum oven for 4 hours6. Package the final, stabilized plates/strips in an airtight container with desiccant <p>To Stabilize Adsorbed or Immobilized Proteins on Membranes</p> <ol style="list-style-type: none">1. Immobilize or adsorb the primary protein (antibody or antigen) according to the procedure optimized in your laboratory2. Dilute 1 part Immunoassay Blocking Buffer stabilizer in three parts compatible buffer or deionized water; add 0.01% surfactant3. Coat membrane by incubating or spraying with the Immunoassay Blocking Buffer stabilizer solution4. Dry thoroughly. Faster drying results in better flow properties5. Package the final, stabilized membrane in an airtight container with a desiccant

To Stabilize Conjugates in the Dry Form

1. If you dilute your conjugate before drying/lyophilizing, use Immunoassay Blocking Buffer stabilizer as the diluent. Otherwise, add between 5-10 parts Immunoassay Blocking Buffer stabilizer or 1 and 2 parts 5X Immunoassay Blocking Buffer stabilizer to 1 part conjugate
2. If you are lyophilizing in vials with rubber stoppers, stability may be improved by placing the rubber stoppers in a 100°C vacuum oven for one hour prior to use; this dries them and drives off any volatiles
3. Freeze the conjugate/stabilizers mixture, then lyophilize as normal; lyophilization may require extra time
4. For evaporation drying, place the conjugate/stabilizer mixture in a 37°C-40°C oven for four hours or until completely dry; the volume per container should be low enough to allow maximum surface area to be exposed to air during drying; store the final product in an airtight container

製品の特徴

製品の状態	Liquid
保存方法	Store at room temperature.
バッファー	pH: 7.00 Constituents: 10% Sucrose, 1% Sodium chloride, 1% Bovine serum protein


アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab171534の使用に適用されず
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ELISA		Use at an assay dependent concentration.

追加情報 Is unsuitable for IHC-Fr or IHC-P.

画像



ELISA - Immunoassay Blocking Buffer (ab171534)

Stability tests were performed with Immunoassay Blocking Buffer (ab171534) using either monoclonal or polyclonal IgG antibodies.

Coated microtiter plates were stabilized with either ab171534 or 1% BSA. The antibody-coated plates were dried and placed at room temperature. Percent retained activity was determined by comparing the activity of the aged antibody-coated microtiter plates to that of freshly coated plates.

After 30 months in Immunoassay Blocking Buffer (ab171534) at room temperature, the monoclonal antibody retained 70% activity and the polyclonal antibody retained 95% activity.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors