

Goat Anti-Rabbit IgG H&L (Alexa Fluor® 405) ab175652

★★★★★ [2 Abreviews](#) [33 References](#) [画像数 4](#)

製品の概要

製品名	Goat Anti-Rabbit IgG H&L (Alexa Fluor® 405)
由来種	Goat
ターゲット生物種	Rabbit
アプリケーション	適用あり: IHC-Fr, ELISA, Flow Cyt, IHC-P, ICC/IF
免疫原	The details of the immunogen for this antibody are not available.
標識	Alexa Fluor® 405. Ex: 402nm, Em: 421nm

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle. Stable for 12 months at -20°C. Store In the Dark.
バッファー	Preservative: 0.02% Sodium azide Constituents: 23% Glycerol (glycerin, glycerine), PBS, 1% BSA
精製度	Immunogen affinity purified
特記事項 (精製)	This antibody was isolated by affinity chromatography using antigen coupled to agarose beads.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG
特記事項	We recommend storage time at 4°C should be minimal, since this may affect the signal strength. Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab175652の使用に適用されます
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-Fr	★★★★★ (1)	Use at an assay dependent concentration.
ELISA		Use at an assay dependent concentration.
Flow Cyt	★★★★★ (1)	1/2000.
IHC-P		Use at an assay dependent concentration.
ICC/IF		1/200 - 1/1000. We recommend the use of a dedicated 405 filter for optimal results not the DAPI filter. The DAPI filter may not excite until the maximum emission peaks of Alexa Fluor® 405 dye (see difference below) Ex max: Alexa Fluor® 405 = 402nm / DAPI = 359nm Em max: Alexa Fluor® 405 = 421nm / DAPI = 461nm

画像



ICC/IF image of **ab6046** stained HeLa cells. The cells were 4% PFA fixed (10 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to block non-specific protein-protein interactions. The cells were then incubated with the antibody (**ab6046**, 5µg/ml) overnight at +4°C. The secondary antibody (blue) was ab175652 Alexa Fluor® 405 goat anti-rabbit IgG (H+L) used at 2µg/ml for 1h. DRAQ5™ (**ab108410**) was used to stain the cell nuclei (red) at a concentration of 1.25µM.




ELISA - Goat Anti-Rabbit IgG H&L (Alexa Fluor® 405) (ab175652)

Cross-reactivity of the polyclonal secondary antibody **ab182016** was tested using a sandwich ELISA approach. The wells were coated with the indicated IgG standards at 1 µg/ml (50 µl/well) and incubated overnight at 4°C, followed by a 5% BSA blocking step for 2h at RT. **ab182016** was then added starting at 1 µg/ml and gradually diluted 1/4 (50 µl/well), followed by incubation for 2h. For the detection Donkey anti-Goat IgG H&L (HRP) (**ab6885**) was used at 1/10,000 dilution (50 µl/well), followed by incubation for 1h at RT.

For the batch tested, ab182016 showed a cross-reactivity of 5-7% towards Human IgG and below 2% towards Mouse IgG, Rat IgG and Chicken IgY.


This data was developed using the unconjugated antibody (**ab182016**).



ELISA - Goat Anti-Rabbit IgG H&L (Alexa Fluor® 405) (ab175652)

Cross-reactivity of Goat anti-Rabbit IgG H&L (**ab182016**) and Goat anti-Rabbit IgG H&L obtained from two different vendors was tested using a sandwich ELISA approach. The wells were coated with the indicated IgG standards (Rabbit, Human, Mouse and Rat) at 1 µg/ml (50 µl/well) and incubated overnight at 4°C, followed by a 5% BSA blocking step for 2h at RT. Secondary antibodies were then added starting at 1 µg/ml and gradually diluted 1/4 (50 µl/well), followed by incubation for 2h. For the detection Donkey anti-Goat IgG H&L (HRP) (**ab6885**) was used at 1/10,000 dilution (50 µl/well), followed by incubation for 1h at RT. This data is from a representative dilution.

This data was developed using the unconjugated antibody (**ab182016**).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors