

### Goat Anti-Rabbit IgG H&L (HRP) preadsorbed ab7090

★★★★★ [2 Abreviews](#) [552 References](#) [画像数 2](#)

#### 製品の概要

製品名	Goat Anti-Rabbit IgG H&L (HRP) preadsorbed
由来種	Goat
ターゲット生物種	Rabbit
アプリケーション	<b>適用あり:</b> ICC/IF, IHC-P, WB, ELISA, Immunomicroscopy, Dot blot, IHC-Fr
吸着処理血清	Chicken, Cow, Goat, Guinea pig, Hamster, Horse, Human, Mouse, Rat, Sheep <a href="#">more details</a>
免疫原	Full length native Rabbit IgG (purified).
標識	HRP

#### 製品の特性


製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	pH: 6.50 Preservative: 0.01% Gentamicin sulphate Constituents: 0.42% Tripotassium orthophosphate, 0.87% Sodium chloride, 1% BSA
精製度	Affinity purified
特記事項 (精製)	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads.
特記事項 (標識)	Horseradish Peroxidase (HRP)
ポリモノ	ポリクローナル
アイソタイプ	IgG

#### アプリケーション

The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab7090の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		1/1000 - 1/5000.
IHC-P	★★★★★ (1)	Use at an assay dependent concentration.
WB	★★★★★ (1)	Use at an assay dependent concentration.
ELISA		1/20000 - 1/80000.
Immunomicroscopy		Use at an assay dependent concentration.
Dot blot		Use at an assay dependent concentration.
IHC-Fr		Use at an assay dependent concentration.


## 画像



Immunohistochemical analysis of healthy and colitis mouse colon sections (untreated and treated with enoxaparin) labeling claudin-4 with **ab15104** at 1/200. ab7090, anti-rabbit immunoglobulin G conjugated to horseradish peroxidase (HRP) at 1/300 was used as the secondary antibody. **ab64238** at 1/50 was used to develop the histological signal.

Antigen retrieval was performed by incubating the sections for 10 minutes at 97°C in 1 mM EDTA buffer, pH 8.0 or 10 mM citrate buffer, pH 6.0.

Control, C; untreated colitis, DSS; oral enoxaparin, OE; intraperitoneal injection of enoxaparin, IPE.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Goat Anti-Rabbit IgG H&L (HRP) preadsorbed (ab7090)

This image is courtesy of an anonymous Abreview.

**ab9986** staining osteoprotegerin in human skin tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 1% BSA and 5% normal goat serum in PBS for 30 minutes at 22°C; Samples were incubated with primary antibody (1/20) for 9 hours at 4°C. An HRP-conjugated goat anti-rabbit IgG H&L (ab7090) (1/300) was used as the secondary antibody.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors